

GÁBOR TAMÁS



University of Szeged
 Albert Szent-Györgyi Medical School
 Department of Physiology, Anatomy and Neuroscience
 MTA-SZTE Research Group for Cortical Microcircuits

Address: Közép fasor 52., H-6726 Szeged, Hungary

RESEARCH AREA

Our research is characterized by a combination of technically challenging electrophysiology, molecular biology, imaging and anatomy in pursuit of the function of cell types and their synapses in the human and rodent cerebral cortex. We discovered the cellular source (neurogliaform cells) of slow, GABAB receptor mediated inhibition in the cerebral cortex. Subsequently, we discovered the mechanism of this slow inhibition as single neuron driven nonsynaptic or volume transmission of the neurotransmitter GABA. In addition, our experiments assigned a new, excitatory role to axo-axonic cells, which were considered as the most specific inhibitory neurons of the cortex. Our commitment to cutting edge methodology recently resulted in recordings from identified interneurons in completely unaesthetized, freely behaving rodents and identified the first ripple-like oscillatory events in the neocortex and their cellular structure. We initiated a research program in 2004 for multiple patch clamp recordings in slices taken from the human cerebral cortex leading to the first recordings of human synaptic interactions and showing the existence of Hebbian networks in the human cerebral cortex.

TECHNIQUES AVAILABLE IN THE LAB

In vivo juxtacellular recordings from neurons of the cerebral cortex in freely behaving rodents, *in vivo* patch clamp electrophysiology, human *in vitro* brain slice patch clamp electrophysiology, *in vivo* and *in vitro* multiphoton imaging (acoustooptical and resonant scanning), CARS microscopy in brain slices, transmission electron microscopy, 3D neuron reconstruction with Neurolucida, single digital PCR, single and oligocellular next generation sequencing.

SELECTED PUBLICATIONS

- Averkin, R., Szemenyei, V., Borde, S., **Tamas, G.** (2016) Identified cellular correlates of neocortical ripple and high-gamma oscillations during spindles of natural sleep. **Neuron** **92**: 916-92.
- Molnar, G., Rozsa, M., Baka, J., Holderith, N., Barzo, P., Nusser, Z., **Tamas, G.** (2016) Human pyramidal to interneuron synapses are mediated by multi-vesicular release and multiple docked vesicles. **eLife** **5**: e18167.
- Olah, S., Fule, M., Komlosi, G., Varga, C., Baldi, R., Barzo, P., **Tamas, G.** (2009) Regulation of cortical microcircuits by unitary GABA-mediated volume transmission. **Nature** **461**: 1278-81.
- Szabadics, J., Varga, C., Molnar, G., Olah, S., Barzo, P., **Tamas, G.** (2006) Excitatory effect of GABAergic axo-axonic cells in cortical microcircuits. **Science** **311**: 233-5.
- Tamas, G.**, Lorincz, A., Simon, A., Szabadics, J. (2003) Identified sources and targets of slow inhibition in the neocortex. **Science** **299**: 1902-1905.